



Browlift and Blepharoplasty Patient Instructions

Pre-operative instructions:

- Please discontinue the following medications two weeks before your planned procedure as they increase the risk of bleeding:
 - Aspirin or aspirin-containing products (Excedrin[®], Alka-Seltzer[®], Pepto-Bismol[®])
 - Ibuprofen (Advil[®], Motrin[®])
 - Naproxen (Aleve[®])
 - Vitamin E or other herbal supplements

You may safely use Tylenol for pain control if needed. Please notify Dr. Friedman's office if you are currently taking warfarin (Coumadin[®]) or clopidogrel (Plavix[®]).

- If you use tobacco products, please refrain from using tobacco products and avoid exposure to secondhand smoke for at least six weeks prior to your procedure as tobacco products decrease the blood flow to the skin and may compromise the healing process.
- Do not eat or drink anything after midnight the night before your procedure.

Post-operative instructions:

Anesthesia Your procedure will most likely be done under local anesthesia with sedation; however, there is a possibility of having it under general anesthesia. Either way, you will need to have someone drive you to and from the hospital on the day of your procedure. If you have general anesthesia, you may feel fatigued for a couple of days following the procedure.

Swelling and bruising After your procedure, you may have moist gauze packed with ice lying over your eyes. This is to decrease the swelling and provide comfort. There will be a moderate amount of swelling and bruising following your procedure. This will be primarily concentrated around the eyes and possibly over the forehead. You should not drive if the swelling is obstructing your vision. You may use ice packs as needed for swelling and bruising, 20 minutes on then 20 minutes off for the first 24-48 hours after surgery. You should attempt to ambulate as early as possible after the procedure since lying in bed can worsen the swelling. You may need assistance when ambulating for the first 24-48 hours after surgery due to swelling. When lying down, it may be helpful to elevate the head of the bed for at least one week. The swelling will subside slowly each day with significant improvement at approximately two weeks post-op. If you had a browlift, you may also experience alterations in your sensation over the forehead and scalp, such as numbness or tingling. This is due to the disruption of nerves during the procedure. This is temporary and should resolve over the next few weeks. Please be careful when using a hair dryer or curling iron as you will not sense the heat due to the numbness over the scalp. You may feel that the brows are elevated more than expected but please remember that they will settle into a more natural position over time.

Wound care All sutures are dissolvable and there is no need for suture removal. If staples are present, they will need to be removed in two weeks. You will be given prescriptions for Bacitracin[®] and Erythromycin[®] ointment to be used on your incision sites. Please apply the Bacitracin[®] ointment to the incisions on your forehead or scalp twice daily for two weeks. The Erythromycin[®] ointment should be applied to the eyelid incisions also twice daily for two weeks. If you should develop crusting over the incision sites, please cleanse gently with soap and water and follow by gently applying half-strength hydrogen peroxide (hydrogen peroxide mixed half-and-half with water) to the area with a Q-tip twice daily. Please take extra care to avoid sun exposure until the wounds are healed and then protect the incision sites with sunscreen as sun exposure can interfere with the healing process. Please refrain from washing your hair for the first 3 days following your procedure if you have had a browlift to prevent disruption of the incision sites.

Vision You may notice that your eyes are more light-sensitive following the procedure and you may wear sunglasses as needed during this time. You may also experience slight dryness of your eyes and should use over-the-counter Artificial Tears[®] as needed. Please refrain from wearing contact lenses until your follow-up appointment in the clinic in approximately two weeks. Also refrain from wearing eye make-up during this time as this can cause "tattooing" of the incision as well as irritation.

Activities Again, please ambulate as soon as possible following your procedure to avoid swelling that is worsened from lying down and to prevent the formation of blood clots. You may return to your daily activities as soon as you feel you are capable, however, you should avoid heavy lifting (greater than 10 lbs) or strenuous exercise for at least two weeks. Although it varies from one individual to the next, you will typically be able to return to work within 1-2 weeks following your procedure.

Medications In addition to prescriptions for Bacitracin® and Erythromycin®, you will also receive prescriptions for narcotic pain medication and an antibiotic. You may use Tylenol® for mild discomfort. Again, please avoid any analgesics that may promote bleeding (see pre-op instructions). If Tylenol® does not adequately control your pain; you may utilize the prescription pain medication. Please note that the prescription pain medication is a Tylenol®-containing product and the maximum dose of Tylenol® is 4 grams (4000mg) in a 24-hour period. You should not drive while taking narcotic pain medication. You should complete the entire course of the antibiotic. Eating yogurt with active cultures may minimize the likelihood of developing diarrhea. If you should develop diarrhea, please stop taking the antibiotic and call Dr. Friedman's office at 215-829-5180 or call the Pennsylvania Hospital operator at 215-829-3000 and ask to be connected to the ENT physician on-call. You may also go directly to the emergency room.

Complications Please monitor for signs of infection including redness, worsening pain and tenderness to touch, an increase in swelling or purulent (pus) drainage from the incision site. Please notify Dr. Friedman's office promptly at 215-829-5180 for any visual changes such as double vision, loss of vision or severe pain. If we are unavailable, please go to the emergency room immediately.

Follow-up appointment You will have a follow-up appointment with Dr. Friedman or Dr. Friedman's assistant approximately two weeks following your procedure.